



# Effectiveness of Child Passenger Safety Information For the Safe Transportation of Children

Kelli England Will, Ph.D,<sup>1</sup> Lawrence E. Decina, MS,<sup>2</sup> Erin L. Maple, MPH,<sup>1</sup> and Amy M. Perkins, MS<sup>1,3</sup>

## Background

Age-appropriate restraints and rear seating dramatically reduce injury in a collision (Arbogast, Jermakian, Kallan, & Durbin, 2009; Durbin, Chen, Smith, Elliott, & Winston, 2005; National Highway Traffic Safety Administration, 2010; Rice & Anderson, 2009). The primary reasons for injuries to children restrained at the time of motor vehicle crashes relate to prematurely turning a child forward, premature graduation from harnessed safety seats to booster seats, premature graduation from booster seats to adult safety belts, misuse of safety restraints and seat belts, and children seated in the front seat of the vehicle (Arbogast et al., 2009; Durbin et al., 2005; Henary et al., 2007; Lennon, Siskind, & Haworth, 2008; Rice & Anderson, 2009). Compared to appropriately restrained children, unrestrained children are 3 times more likely to sustain injury in a crash, and children traveling in *inappropriate* restraints for their size are at twice the risk of injury (Durbin et al., 2005). Rear seating offers independent and additive safety protections in a crash (Durbin et al., 2005; Lennon et al., 2008).

A large number of studies over the past decade have involved some type of intervention to increase the correct use of child restraints, including booster seats and seat belts. Most of these efforts included educational material and messages as part of the interventions (Dellinger, Sleet, Shults, & Rinehart, 2007; Dukehart, Walker, Lococo, Decina, & Staplin, 2007; Ebel, Koepsell, Bennett, & Rivara, 2003; King, Monroe, Applegate, & Cole-Farmer, 2007; Snowdon et al., 2008; Weiss-Laxer, Mello, & Nolan, 2009; Winston, Erkoboni, & Xie, 2007; Zaza, Sleet, Thompson, Sosin, & Bolen, 2001). However, very few of these studies dealt specifically with evaluation of the messaging

associated with these interventions. For those studies that looked at messaging, research suggests that messages that increase parents' feelings of vulnerability to risk and provide succinct and concrete educational messages about the injury prevention benefits of car seats will be most likely to increase correct use of child restraints for children (Sheeran, Harris, & Epton, 2014; Will, 2005; Will, Sabo, & Porter, 2009; Winston et al., 2007). Research also indicates it is important to depict negative consequences in parental safety messages in order to effectively communicate danger and evoke attention and concern (Morrongiello, Bell, Butac, & Kane, 2013). Combating parents' low perceptions of risk for motor vehicle injury will likely be difficult since the risk of being involved in a crash on any given vehicle trip is very small, which in turn reinforces the perception of minimal risk (Will, 2005; Will & Geller, 2004).

## Project Objectives

The objective of this project was to develop and test various methods of framing child passenger safety recommendations for children under age 13. This research note reports on the first of two studies examining child passenger safety messages and types of information on which to focus for maximum effectiveness. The goal of this first study was to determine how to best communicate child passenger safety recommendations to parents/caregivers, and which information to emphasize. Thus, this study investigated various ways of framing child passenger safety recommendations, and examined the relative effectiveness on parents/caregivers' knowledge, attitudes, and behavioral intentions related to best practices and proper use of child restraints. Specifically, should the recommendations be organized by phase of childhood (e.g., by age, or by progression of younger to older)? Should they focus on key issues, such as combating premature graduation? Should they communicate risk-reduction rationale and consequences of noncompliance? Note that the base child passenger safety recommendations are consistent across

<sup>1</sup>Eastern Virginia Medical School, Norfolk, VA

<sup>2</sup>TransAnalytics, Quakertown, PA

<sup>3</sup>Children's Hospital of The King's Daughters, Norfolk, VA

conditions in this study, but several versions are tested that each employ a different emphasis frame. Emphasis framing is a persuasion technique that involves placing focus on specific aspects of the content in order to encourage or discourage certain interpretations of the content. Considerable research indicates that varying communication frames can affect attitudes and behaviors, even among two otherwise equivalent statements (Chaiken, 1987; Chong & Druckman, 2007; Kahneman, Slovic, & Tversky, 1982).

It was hypothesized that the varying emphasis frames would have a differential effect on knowledge, attitudes, and behavioral intentions, despite the base CPS recommendations being consistent across conditions. Further, it was hypothesized that all experimental frames would be more effective than the materials viewed in the control condition, and the frame that explained the risk-reduction rationale behind the recommendations would be most effective at improving knowledge, attitudes, and behavioral intentions.

## Study Methods

### *Study Design*

A 5 (test conditions) × 2 (time periods) experiment was conducted using a randomized controlled trial design to examine relative effectiveness of parent and caregiver preferences for different methods of framing car seat safety recommendations. Participants were electronically randomized to 1 of 5 test condition groups (4 experimental conditions and 1 control group) and responded to pre- and post-survey questions (2 times).

### *Sampling Plan*

The study took place in the suburbs of Philadelphia, Pennsylvania, and in Norfolk/Hampton Roads, Virginia. These socio-economically and culturally diverse areas covered urban and suburban concentrations of candidate parents and caregivers. Each site recruited and tested 150 participants each (300 total sample).

### *Recruitment and Incentives*

Each site used various methods to advertise the study to parents or caregivers of children from birth to 12 years old. For the Philadelphia site, the team worked with the Safe Kids Chapter of Southeast Pennsylvania to deliver recruitment flyers to various parent clubs, online parent newspapers, and child-care facility organizations. For the Norfolk site, the team used very similar methods, working with child-focused organizations (e.g., Places and Programs for Children, Consortium for Infant and Child Health,

etc.) to deliver the recruitment flyers to various groups of parents and child care facilities through their contact networks. Facebook sites were also used to promote the study at both sites. Scheduling of participants was handled through e-mail communications and telephone correspondence. Each site had various days and times set up for parents/caregivers to participate in the study in a local computer lab setting. Participants were compensated with a \$50 Walmart gift card for their participation in the study.

### *Procedures for Participation*

Enrolled participants were asked to arrive at a designated computer lab center at their appointment time to participate in the study. A secure Web-based study protocol was used for participants who viewed a series of user-friendly screens that automatically led them through an informed consent document (covering logistics of study, duration, rights as a participant, and remuneration for participation), pretest measures, study material specific to condition assignment, and post-test measures, at their own pace. Most participants were able to complete the session in about one-half hour (mean = 26 minutes). A study facilitator was present at all sessions to assist the participants in log-in procedures, to answer any questions and to resolve any administrative issues. The sessions were also monitored off-site by the Web site designer to confirm data recording. No person's name or other personal identifiers were stored with the data; an anonymous coding process was used to link pre- and post-data. Upon completion of the testing session, participants signed for and received their compensation, as well as a handout on child passenger safety to take home for reference.

### *Test Study Conditions*

Participants were electronically randomized to one of the five test groups to view child passenger safety material. Four groups viewed one of four versions of a one-page educational print flyer, and one control group viewed car seat marketing material that were not educational. Randomizing participants to groups allowed for examination of the relative effectiveness of and preferences for different methods of framing child restraint recommendations. The five groups are described below.

#### Group 1: Child Restraint Recommendations Organized by Natural Progression

Participants in this group viewed a version of car seat recommendations in which both text and pictures highlighted the natural progression of seat types from birth to teen years (see Figure 1). This version used photos of children representative of each phase of childhood, but removed almost all references to age and all mention of

upper limits for common seats as a factor for determining transitions. This message frame was chosen to examine the utility of organizing CPS recommendations by phase of childhood (younger to older progression) without attaching specific age ranges to the phases. Thus, this version was the most similar to the “Organized by Age” flyer (see Group 4 below), but does not include references to ages in the organizing headers. Recommendations for transitioning from rear-facing to forward-facing pushed toward later transition. To quell the perception that age 8 is the maximum, it is mentioned that it may take up to 12 years old for a child to be big enough to use a seat belt alone. Recommendations for this condition focused on best practice for determining transitions to the next stage, which include child size and fit of the restraint. For instance, transition to seat belts focused on fit of the belt on the seated child (using the fit test), with usual maximum height for a booster seat (4’ 9”) given as additional guides. Pictures of older children for each phase were used to emphasize the upper transition norms for each stage. The need for back seat positioning was fully integrated and highlighted throughout the recommendations.

#### Group 2: Child Restraint Recommendations That Focus on Premature Graduation

Participants in this group viewed a version of car seat recommendations in which both text and pictures draw attention to premature graduation (see Figure 2). This message frame was chosen to determine the value of organizing CPS information around the best practice guidance for delaying transitions between stages of child restraint. In addition to specifying recommendations for each stage, this version specifically emphasized the message that counters premature graduation to the next stage. For instance, the header for stage two read, “Keep Kids in Seats with Harnesses as Long as Possible” to emphasize the need to use harnesses throughout this stage. Parents were encouraged to keep children in harnessed seats for as long as the harness weight and height limits will allow. Similar encouragements against premature graduation were used for each phase as was appropriate for the phase. Accompanying pictures provided additional emphasis. Similar to Group 1, this version also removed almost all references to age and upper limits for common seats, and fully integrates and highlights the need for back seat positioning at all stages.

#### Group 3: Child Restraint Recommendations That Explain Risk-Reduction Rationale

Participants in this group viewed a version of car seat recommendations that communicated the risk-reduction potential and rationale (in a lay-friendly, succinct manner avoiding statistics) behind each stage’s recommended

restraint configuration, starting first with the basic rationale for occupant restraints and moving into rationale for specific restraint configurations for the various child sizes. This message frame (see Figure 3) is consistent with risk communication literature for maximum behavior change, and was chosen to examine the merits of focusing on *why* each seat/configuration makes a difference for safety. Much of the general public fails to recognize the severity of many public health hazards, including motor vehicle travel (Sandman, 1989; Slovic, 1991; Slovic, Fischhoff, & Lichtenstein, 1985; Will & Geller, 2004). For instance, many parents may lack the understanding that an object in motion remains in motion, unless restrained, when the vehicle crashes. They may also fail to grasp that given the abrupt changes in momentum and velocity that occur in mere fractions of a second, crash forces are quite powerful and can result in a child propelling forward with the force of thousands of pounds (National Child Passenger Safety Board, 2014). Thus, the *reasons* behind the recommendations were given for each stage in simple “Here’s What to Do” and “Here’s Why” sections. For instance, parents were not only told to rear-face their children longer, but why rear-facing provides such a benefit in crashes. This version also included pictures to illustrate stages of restraints, removed almost all references to age and upper limits for common seats, and fully integrated the need for back seat positioning at all stages.

#### Group 4: Child Restraint Recommendations Organized by Age

Participants in this group viewed a version of car seat recommendations that are organized under age-based headers (see Figure 4). An age-based frame was included for examination, given its frequent use in CPS and health-related communications to parents. The National Highway Traffic Safety Administration’s “Car Seat Recommendations for Children” flyer (released in March 2011) was used as the flyer for this group. The flyer focuses on age of child for specific type of car seat or restraint; and fit of child based on car seat manufacturer’s instructions for size and height of child. The flyer emphasizes importance of harnesses and seat belt positions for rear-facing and forward-facing car seats, as well as booster seats and seat belts. The flyer mentions the need to read the vehicle owner’s manual on how to install the car seat using the seat belt or Lower Anchors and Tethers for Children (LATCH) system, and the need to check height and weight limits.

#### Group 5 (Control): No Education

Participants assigned to this condition did not receive any instructional material related to car seats. Rather, these participants viewed a picture display of various car seats on the market and were asked to rate their preferences based on style, color, and other characteristics. This exer-

cise allowed for elapsed time between their pretest and post-test measures, as in the other study conditions, without providing education.

### Measures

Several measurement scales were used to measure appropriateness of restraint selection, knowledge of restraints, perceived efficacy and threat, attitudes and intentions, judgments of relevance and acceptability, and sample demographics. To accurately assess changes in knowledge and perceptions after exposure to the independent variable, most of the measures were asked in both the pretest and post-test. The exceptions were the demographics questions and judgments of relevance and acceptability, which were asked only once (at post-test).

### Restraint Selection

For proper child restraint selection, items were developed that provided participants with a series of specific scenarios that vary the age, weight, and height of a child and asks them to select an appropriate restraint, direction to face, and vehicle row for the hypothetical child. This 8-item knowledge measure used a multiple choice response format, providing an item score of correct/incorrect and a total number correct score for each participant. A sample question is, "Your child is asking you when he/she can just use a seat belt when riding in vehicles. When can your child safely use a seat belt only?" The measure was adapted from a similar existing field-tested measure (Snowdon et al., 2008).

### Child Passenger Safety Knowledge

To gauge immediate changes in general knowledge of child passenger safety, as well as differences in knowledge among the groups, a 15-item assessment of parental knowledge was conducted at both pretest and post-test. This assessment included separate subscales for Back Seat Knowledge, Booster Seat Knowledge, Rear-Facing Knowledge, Forward-facing Knowledge, and Seat Belt Knowledge. This measure used a Likert-type response format (1 = strongly disagree to 5 = strongly agree) and was tailored for this study from existing field-tested and validated measures used in past research by Snowden and colleagues (Snowdon et al., 2008; Snowden, Hussein, Purc-Stevenson, Follo, & Ahmed, 2009), and Will and colleagues (Will et al., 2009). A sample item is, "It is safe for an 11-year-old to ride in the front seat."

### Perceptions of Efficacy and Threat

The Risk Behavior Diagnosis Scale (RBDS) (Witte, Cameron, McKeon, & Berkowitz, 1996) was used to assess perceived efficacy and risk. The RBDS is a template survey

designed to be tailored for evaluation of any health or safety message. The efficacy subscale assessed participants' perceptions of response efficacy (i.e., confidence that the recommended actions/restraints will work to prevent injuries) and self-efficacy (i.e., confidence in one's ability to follow child passenger safety recommendations). The threat subscale assessed participants' perceived risk by measuring susceptibility to and severity of negative consequences from inappropriate child occupant protection. The 16-item RBDS tailored for this study used a 5-point Likert-type response scale (1 = strongly disagree to 5 = strongly agree). A sample self-efficacy item is, "I have the skills and knowledge needed to use the correct restraint to reduce my child's chances of injury in a car crash."

### Attitudes and Intentions

Participant's general attitudes and intentions regarding child passenger safety were assessed via an 8-item attitudes subscale, adapted from a survey used in past research (Will et al., 2009), and a 9-item stated intentions subscale. Stated intentions and attitudes were assessed to gauge participants' disposition regarding what is recommended for child occupant protection irrespective of their knowledge. Both subscales used a Likert-type response format (1 = strongly disagree to 5 = strongly agree). A sample attitudes item is, "Rear-facing a child past the first birthday seems harmful because there is not enough room for his/her legs."

### Judgments of Relevance and Acceptability

At post-test, participants in each experimental condition were asked their opinions about the child passenger safety materials. A 10-item questionnaire was developed that uses a 4-point Likert-type response format to assess participants' judgments of quality and acceptability of the information presented. Specifically, they were asked to rate the child passenger safety information on a variety of factors, including but not limited to style, amount of information, clarity, and likelihood for motivating behavior change. A sample question is, "How would you rate the clarity of the materials that were presented to you today?"

### Demographics and other Participant Information

Demographic information was collected at the post-test. Participants were asked their age, gender, race, ethnicity, education level, income level, and number of and ages of children. Information specific to child passenger safety was also asked, including types of child restraints being used currently and in the past, sources of information about safely transporting children, whether or not they have had their children's restraints inspected by a CPS technician, and their preferred communication channels

(e.g., print, television, radio, electronic) for receiving child passenger safety information.

## Results

### Participants

Three hundred parents/caregivers of children from birth to 12 participated in the study (150 in the suburbs of Philadelphia and 150 in the Norfolk/Hampton Roads area). Demographics are presented in Table 1. Thirty-four percent of participants were Black (59% of the Hampton Roads participants and 9% of the Philadelphia participants). About 3 percent of Philadelphia and 7 percent of Hampton Roads participants were Hispanic. Mean parent age was 38 and 33 years old in Philadelphia and Hampton Roads, respectively. Males made up 11 percent of the participants (13% and 9% for Philadelphia and Hampton Roads areas, respectively). Participants at the two sites did not significantly differ on responses to surveys/flyer versions.

### Analysis

Analyses of covariance (ANCOVA) and pair-wise comparisons with Sidak's adjustment for Type 1 error were used to determine the relationship between group assignment (1 of 4 flyer versions or control) and post-test scores after adjusting for pre-test scores and the interaction between the independent factors when significant. This analysis compares differences among post-scores while controlling for pre-scores. Data were analyzed using SPSS 19 software and the level of significance was set at 0.05. After adjusting for pre-test scores, post-test score means revealed a significant main effect for flyer version on 11 subscales: (a) Restraint Selection Score; (b) Back Seat Knowledge; (c) Booster Knowledge; (d) Rear-Facing Knowledge; (e) Total Efficacy; (f) Self-Efficacy; (g) Overall Attitudes; (h) Booster Attitudes; (i) Forward-Facing Attitudes; (j) Rear-Facing Attitudes; and (k) Stated Intentions. These findings are described separately below.

### Changes in Restraint Selection Score

Analyses for Restraint Selection Score revealed a significant main effect for flyer,  $F(4, 293) = 7.72, p < .001, \eta_p^2 = .10$ , and a significant interaction with pretest score,  $F(1, 293) = 9.07, p = .003, \eta_p^2 = .03$ . Given the significant interaction between pre-test score and flyer version, the effectiveness of each flyer on the post-test score was dependent upon the pre-test score. After analyzing the interaction, post-test scores for the Risk Reduction Rationale flyer were significantly higher than the Age-Based flyer across all pre-test scores ( $p = .02$ ). Additionally, participants viewing the Risk Reduction Rationale or Natural Progression versions performed significantly better than participants in the Control group when pretest scores were below 50 percent

( $p = .003$ ). The Premature Graduation, Age-Based, and Control groups did not differ significantly from one another. Figure 5 depicts mean changes in restraint selection scores by group. In summary, when faced with the task of selecting appropriate restraints for given children, the flyer that provided the rationale behind the recommendations led to the greatest improvement in scores from pre- to post-test.

### Changes in Child Passenger Safety Knowledge

Child passenger safety knowledge analyses revealed a significant main effect for flyer for Back Seat Knowledge,  $F(4, 294) = 2.84, p = .03, \eta_p^2 = .04$ . Specifically, after adjusting for pretest scores, post-test scores for the Risk Reduction Rationale ( $p = .04$ ) and Premature Graduation ( $p = .04$ ) flyers were significantly higher than the control group. For Booster Seat Knowledge, there was a significant main effect for flyer,  $F(4, 293) = 2.59, p = .04, \eta_p^2 = .03$ . While it appears the Premature Graduation version resulted in greater change in booster knowledge, there was an interaction indicating flyers performed differently for high vs. low pre-scores,  $F(1, 293) = 7.11, p = .01, \eta_p^2 = .02$ . Further comparisons for booster knowledge lacked power to reach significance. Regarding Rear-Facing Knowledge, there was a significant main effect for flyer,  $F(4, 293) = 5.33, p < .001, \eta_p^2 = .07$ , and a significant interaction between pre-test score and flyer version,  $F(1, 293) = 9.61, p = .002, \eta_p^2 = .03$ . After analyzing the interaction, participants viewing the Risk Reduction Rationale ( $p = .03$ ) or Natural Progression ( $p = .003$ ) versions performed significantly better than participants in the Control group for all but the highest 20 percent of pretest scores. Participants in the Premature Graduation version performed somewhere in the middle, no better than those in the Control and no worse than those in the Risk Reduction Rationale and Natural Progression groups. Figure 6 depicts mean changes in rear-facing knowledge by group. No significant main effects for flyer were evident for Seat Belt Knowledge or Forward-facing Knowledge subscales. In summary, the greatest changes in knowledge were found most often among participants who viewed either the flyer that emphasized the rationale behind the recommendations, or the one that focused on dissuading premature graduation.

### Changes in Perceptions of Efficacy and Threat

Analyses of threat and efficacy subscales revealed a significant main effect for flyer for Total Efficacy,  $F(4, 294) = 3.64, p = .01, \eta_p^2 = .05$ , and for Self-Efficacy,  $F(4, 294) = 3.48, p = .01, \eta_p^2 = .05$ . After adjusting for pretest scores, participants viewing the Risk Reduction Rationale ( $p = .02$ ) or Premature Graduation ( $p = .01$ ) flyers reported significantly higher total efficacy than participants in the Control group at post-test. Participants viewing the

Natural Progression or Age-Based versions did not differ significantly from the Control group. After adjusting for pretest scores, participants viewing the Premature Graduation flyer had significantly higher self-efficacy scores at post-test ( $p = .01$ ) compared to those in the Control group. No other groups differed from Control for self-efficacy. Figure 7 presents changes in self-efficacy scores by group. No significant main effects for flyer were evident for the Threat Perceptions (Severity and Susceptibility) or Response Efficacy Subscales. In summary, participants who viewed the flyer that focused on dissuading premature graduation exhibited greater increases in efficacy compared to participants in other groups.

### Changes in Attitudes and Intentions

Analyses of the Overall Attitudes scale revealed a significant main effect for flyer,  $F(4, 294) = 8.03, p < .001, \eta_p^2 = .10$ . After adjusting for pretest scores, the Natural Progression ( $p < .001$ ), Premature Graduation ( $p < .001$ ), Risk Reduction Rationale ( $p < .001$ ), and Age-Based ( $p = .01$ ) flyers performed significantly better than the Control condition in changing attitudes, but the four flyers did not significantly differ from one another. Figure 8 presents mean changes in overall attitudes by group. Regarding the Booster Attitudes subscale, there was a significant main effect for flyer,  $F(4, 294) = 5.00, p = .001, \eta_p^2 = .06$ , where post-test scores for the Risk Reduction Rationale ( $p = .01$ ), Natural Progression ( $p = .001$ ), and Premature Graduation ( $p = .004$ ) flyers were significantly higher than Control after adjusting for pretest scores. There was also a significant main effect for flyer for the Forward Facing Attitudes subscale,  $F(4, 294) = 2.51, p = .04, \eta_p^2 = .03$ . However, pairwise comparisons lacked power to reach significance. It appears that all 4 flyers were likely different from control in their ability to change forward-facing attitudes. Analyses for the Rear-facing Attitudes subscale revealed a significant main effect for flyer,  $F(4, 294) = 6.43, p < .001, \eta_p^2 = .08$ . Pairwise comparisons for Rear-facing Attitudes revealed post-test scores for the Risk Reduction Rationale ( $p = .001$ ), Natural Progression ( $p = .001$ ), and Premature Graduation ( $p = .002$ ) flyers were all significantly higher than the Control after adjusting for pretest scores. No significant main effect for flyer was evident for the Back Seat Attitudes subscale. In summary, all four flyer versions performed equally well in regards to overall attitude change; however, the Age-Based flyer did not differ from Control when looking specifically at booster attitudes and rear-facing attitudes.

Analyses for parents' Stated Intentions revealed a significant main effect for flyer,  $F(4, 293) = 5.00, p = .001, \eta_p^2 = .06$ , and a significant interaction with pretest score,  $F(1, 293) = 12.87, p < .001, \eta_p^2 = .04$ . Given the significant

interaction between pre-test score and flyer version, the effectiveness of each flyer on the post-test score was dependent upon the pre-test score. For all except the highest pre-scores, Risk Reduction Rationale ( $p = .04$ ), Natural Progression ( $p = .04$ ), and Age-Based ( $p = .04$ ) flyers all resulted in significantly higher post-scores than control. Additionally, except for the highest pre-scores, the Risk Reduction Rationale flyer resulted in significantly higher post-scores than the Premature Graduation flyer ( $p = .04$ ). Figure 9 depicts mean changes in stated intentions by group.

### Judgments of Relevance and Acceptability of Material

Fisher's exact test was used to examine differences among flyers in parents' perceptions of the materials presented. All flyers were rated favorably, and no significant differences were found among the four informational flyers for any of the subscales. Regarding parents' perceptions of the materials presented, 99.2 percent of parents found the information to be organized and coherent, 99.6 percent reported the material to be relevant to their needs, and 97.9 percent believed the amount of information given was appropriate. Regarding parents' perceptions of the quality and clarity of materials, 98 percent found the quality of materials to be good or excellent, and 96.3 percent reported the clarity of the information to be good or excellent. Regarding parents' perceptions of learning and likelihood of changing behavior, 88 percent reported that they gained some or a lot of knowledge, and 83.1 percent believed they were likely to change behavior as a result of receiving the information.

### Conclusions

The Risk Reduction Rationale flyer outperformed other flyers for many subscales, and significantly differed from control for the most subscales, including restraint selection, back seat knowledge, rear-facing knowledge and attitudes, total efficacy, overall attitudes, and stated intentions. The Premature Graduation flyer performed best for efficacy subscales, but did not significantly differ from the Risk Reduction Rationale flyer for total efficacy. For changes in self-efficacy, the Premature Graduation flyer outperformed all other flyers. The Natural Progression flyer performed best for attitudes subscales, but did not significantly differ from the Risk Reduction Rationale flyer. The Age-Based flyer performed significantly better than control only for changes in overall attitudes and stated intentions. However, the Age-Based flyer was outperformed by changes produced by the Risk Reduction Rationale flyer for restraint selection score. All materials were rated favorably, with no significant differences among flyers for parent's ratings.

It is important to note that although the Risk Reduction Rationale flyer was the most effective in a laboratory setting, the flyer has a great deal of information on it and thus needs work in order to be ready for the market. What this study has revealed is that the most advantageous way of framing CPS recommendations is to explain the injury risks behind the information given. The challenge that remains is for marketers and communications professionals to figure out how to present injury risks in a way that is aesthetically pleasing and reader-friendly.

## Recommendations

The objective of this project was to develop and test various methods of framing child passenger safety recommendations, and to determine how to best communicate child passenger safety information to parents/caregivers, and which information to emphasize. A number of recommendations are evident from this research.

1. *Communicate risk-reduction rationale behind the recommendations.* With its focus on the reasons underlying the recommendations, the Risk Reduction Rationale flyer outperformed all other flyers on the most subscales. It is important to tell parents what to do and why it is safer, in straightforward and simple terms. Educational material should communicate the reason behind the recommendation, avoiding use of statistics or abstract comparisons to explain statistics.
2. *Use clear behavior-based directives in headers.* With its action-oriented headers, the Premature Graduation flyer also performed well on a number of subscales, and outperformed all others in its ability to bolster parents' self-efficacy for carrying out the recommendations. Headers should state clearly what needs to be done (e.g., keep kids rear-facing as long as possible), with subtext giving additional details for clarity. It is best to avoid questions in headers and taglines, as it is a missed opportunity for education.
3. *Avoid age-based headers.* In contrast to the Age-Based flyer, the other three flyers did not use age in the organizing headers and to the extent possible, avoided refer-

ences to age and upper limits for common seats within the text. Age is just one of many factors to consider when choosing a restraint. It is known from research on judgmental heuristics that people employ mental shortcuts in making everyday judgments (Chaiken, 1987; Cialdini, 1993; Kahneman et al., 1982). Thus, once an age-based header is used, human nature dictates that many parents will not read the subtext below the header that qualifies the age range given. While it is important to organize the information according to child maturity for parents' ease of use, it is best to convey age progression through the use of stages, arrows, representative pictures, or other means. Information on age parameters can be included in the subtext, but it is not recommended as an organizing header.

4. *Fully integrate the need for back seat positioning at all stages.* Because rear seating offers independent and additive safety protections in a crash, back seat positioning should be recommended concurrently with each restraint configuration. The flyers that fully integrated the need for back seat positioning with each seat recommendation (e.g., "use a booster seat in the back seat") were the most effective in the study.

## For More Information

This Research Note was written by Kelli England Will, an associate professor at Eastern Virginia Medical School (EVMS) in Norfolk; Lawrence E. Decina, a senior associate at TransAnalytics, LLC in Quakertown, PA; Erin L. Maple, a research associate at EVMS, and Amy M. Perkins, a biostatistician employed jointly at EVMS and the Children's Hospital of The King's Daughters in Norfolk. Additional data and information will be available in upcoming reports and publications. For questions regarding the information presented in this document, please contact willke@evms.edu. This research was funded by the National Highway Traffic Safety Administration under contract number DTNH22-09-D-00135.

The suggested APA format for this document is:

Will, K. E., Decina, L. E., Maple, E. L., & Perkins, A. M. (2015, February). *Effectiveness of child passenger safety information for the safe transportation of children*. (Traffic Safety Facts Research Note. Report No. DOT HS 812 121). Washington, DC: National Highway Traffic Safety Administration.

## References

- Arbogast, K., Jermakian, J. S., Kallan, M. J., & Durbin, D. R. (2009). Effectiveness of belt positioning booster seats: An updated assessment. *Pediatrics*, *124*(5), 1281–1286.
- Chaiken, S. (1987). The heuristic model of persuasion. In M. P. Zanna, J. M. Olson & C. P. Herman (Eds.), *Social Influence: The Ontario Symposium* (Vol. 5). Hillsdale, NJ: Lawrence Erlbaum.
- Chong, D., & Druckman, J. N. (2007). Framing theory. *Annual Review of Political Science*, *10*, 103–126.
- Cialdini, R. (1993). *Influence: Science and practice* (3rd ed.). New York: HarperCollins College Publishers.
- Dellinger, A., Sleet, D., Shults, R. A., & Rinehart, C. (2007). Interventions to prevent motor vehicle injuries. In L. Doll, S. Bonzo, D. Sleet, J. Mercy & E. Haas (Eds.), *Handbook of Injury and Violence Prevention*. Atlanta: Springer.
- Dukehart, J. G., Walker, L., Lococo, K. H., Decina, L. E., & Staplin, L. (2007). *Safe Kids checkup events: A national study*. Washington, D.C.: Safe Kids Worldwide.
- Durbin, D., Chen, I., Smith, R., Elliott, M., & Winston, F. (2005). Effects of seating position and appropriate restraint use on the risk of injury to children in motor vehicle crashes. *Pediatrics*, *115*(3), e305–e309.
- Ebel, B. E., Koepsell, T. D., Bennett, E. E., & Rivara, F. P. (2003). Use of child booster seats in motor vehicles following a community campaign. *Journal of the American Medical Association*, *289*(7), 879–884.
- Henary, B., Sherwood, C. P., Crandall, J. R., Kent, R. W., Vaca, F. E., Arbogast, K. B., et al. (2007). Car safety seats for children: Rear facing for best protection. *Injury Prevention*, *13*, 398–402.
- Kahneman, D., Slovic, P., & Tversky, A. (Eds.). (1982). *Judgment under uncertainty: Heuristics and biases*. New York: Cambridge University Press.
- King, W. D., Monroe, K., Applegate, J., & Cole-Farmer, J. (2007). The impact of education, legislation and service on Alabama child passenger safety. *Journal of Trauma*, *63*(3), S25–S28.
- Lennon, A., Siskind, V., & Haworth, N. (2008). Rear seat safer: Seating position, restraint use and injuries in children in traffic crashes in Victoria, Australia. *Accident Analysis and Prevention*, *40*, 829–834.
- Morrongiello, B. A., Bell, M., Butac, M., & Kane, A. (2013). What features of images affect parents' appraisal of safety messages? Examining images from the A Million Messages programme in Canada. *Injury Prevention*, *10.1136/injuryprev-2012-040721*.
- National Child Passenger Safety Board. (2014). *National Child Passenger Safety Certification Training Program technician guide*. Washington, DC: Author.
- National Highway Traffic Safety Administration. (2010). *Traffic Safety Facts 2009: Children* (Report No. DOT HS 811 387). Washington, DC: Author.
- Rice, T. M., & Anderson, C. L. (2009). The effectiveness of child restraint systems for children aged 3 years or younger during motor vehicle collisions: 1996 to 2005. *American Journal of Public Health*, *99*(2), 252–257.
- Sandman, P. M. (1989). Hazard versus outrage in the public perception of risk. In V. T. Covello, D. B. McCallum & M. T. Pavlova (Eds.), *Effective risk communication* (pp. 45–49). New York, NY: Plenum Press.
- Sheeran, P., Harris, P. R., & Epton, T. (2014). Does heightening risk appraisal change people's intentions and behavior? A meta-analysis of experimental studies. *Psychological Bulletin*, *140*(2), 511–543.
- Slovic, P. (1991). Beyond numbers: A broader perspective on risk perception and risk communication. In D. G. Mayo & R. D. Hollander (Eds.), *Acceptable evidence: Science and values in risk management* (pp. 48–65). New York, NY: Oxford University Press.
- Slovic, P., Fischhoff, B., & Lichtenstein, S. (1985). Regulation of risk: A psychological perspective. In R. G. Noll (Ed.), *Regulatory policy and the social sciences* (pp. 241–283). Berkeley, CA: University of California Press.
- Snowdon, A. W., Hussein, A., High, L., Stamler, L., Millar-Polgar, J., Patrick, L., et al. (2008). The effectiveness of a multimedia intervention on parents' knowledge and use of vehicle safety systems for children. *Journal of Pediatric Nursing*, *23*(2), 126–139.
- Snowdon, A. W., Hussein, A., Purc-Stevenson, R., Follo, G., & Ahmed, E. (2009). A longitudinal study of the effectiveness of a multi-media intervention on parents' knowledge and use of vehicle safety systems for children. *Accident Analysis & Prevention*, *41*(3), 498–505.
- Weiss-Laxer, N. S., Mello, M. J., & Nolan, P. A. (2009). Evaluating the educational component of a hospital-based child passenger safety program. *Journal of Trauma-Injury Infection and Critical Care*, *67*(1), S30–S33.
- Will, K. E. (2005). Child passenger safety and the immunity fallacy: Why what we are doing is not working. *Accident Analysis and Prevention*, *37*, 947–955.
- Will, K. E., & Geller, E. S. (2004). Increasing the safety of children's vehicle travel: From effective risk communication to behavior change. *Journal of Safety Research*, *35*, 263–274.
- Will, K. E., Sabo, C. S., & Porter, B. E. (2009). Evaluation of the Boost 'em in the Back Seat program: Using fear and efficacy to increase booster seat use. *Accident Analysis and Prevention*, *41*, 57–65.
- Winston, F. K., Erkoboni, D., & Xie, D. (2007). Identifying interventions that promote belt-positioning booster seat use for parents with low educational attainment. *Journal of Trauma*, *63*(3 Suppl), S29–38.
- Witte, K., Cameron, K., McKeon, J., & Berkowitz, J. (1996). Predicting risk behaviors: Development and validation of a diagnostic scale. *Journal of Health Communication*, *1*, 317–341.
- Zaza, S., Sleet, D. A., Thompson, R. S., Sosin, D. M., & Bolen, J. C. (2001). Reviews of evidence regarding interventions to increase use of child safety seats. *American Journal of Preventive Medicine*, *21*(4 Suppl), 31–47.

Table 1.  
**Demographics of Study Participants by Location**

	Philadelphia Suburb, PA	Hampton Roads, VA	Overall
Number of subjects	150	150	300
Age of parent, mean (SD)	38.1 (5.4)	33.1 (8.1)	35.6 (7.3)
<b>Race, n (%)</b>			
American Indian or Alaska Native	0 (0.0)	0 (0.0)	0 (0.0)
Asian	5 (3.3)	4 (2.7)	9 (3.0)
Black or African American	14 (9.3)	89 (59.3)	103 (34.3)
Native Hawaiian or Pacific Islander	1 (0.7)	1 (0.7)	2 (0.7)
White	129 (86.0)	53 (35.3)	182 (60.7)
Other	1 (0.7)	3 (2.0)	4 (1.3)
<b>Ethnicity, n (%)</b>			
Hispanic	5 (3.3)	10 (6.7)	15 (5.0)
Non-Hispanic	15 (96.7)	140 (93.3)	285 (95.0)
<b>Education, n (%)</b>			
Grade school	0 (0.0)	0 (0.0)	0 (0.0)
Some high school	1 (0.7)	2 (1.3)	3 (1.0)
High school diploma/GED	6 (4.0)	23 (15.3)	29 (9.7)
Some college	15 (10.0)	40 (26.8)	55 (18.3)
2-year degree/trade school	7 (4.7)	32 (21.3)	39 (13.0)
Bachelor's degree	73 (48.6)	38 (25.3)	111 (37.0)
Graduate degree	48 (32.0)	15 (10.0)	63 (21.0)
<b>Income, n (%)</b>			
\$0–15,999	0 (0.0)	18 (12.0)	18 (6.0)
\$16,000–24,999	0 (0.0)	18 (12.0)	18 (6.0)
\$25,000–49,999	16 (10.7)	57 (38.0)	73 (24.4)
\$50,000–99,999	63 (42.0)	37 (24.7)	100 (33.3)
\$100,000+	58 (38.6)	11 (7.3)	69 (23.0)
Decline to answer	13 (8.7)	9 (6.0)	22 (7.3)
<b>Age of children, n (%)</b>			
0–12 months	5 (1.9)	24 (9.3)	29 (5.5)
1–3 years	72 (26.7)	77 (29.8)	149 (28.2)
4–7 years	129 (47.7)	76 (29.5)	205 (38.8)
8–12 years	64 (23.7)	81 (31.4)	145 (27.5)

Figure 1. Natural Progression flyer viewed by participants in Group 1

## Right Seat • Right Time • Right Use

Every seat is different. Every child is different. Here is the natural progression of restraint types from birth to teen years...with a bit of advice for best protection at each stage. Follow seat directions and use each type for as long as possible to the top weight and height limits before transitioning to the next seat type. Unrestrained children are 3 times more likely to be injured. Kids are always safest in the BACK SEAT!

Stage of Childhood	Seat Type	Advice for Stage & Seat Transitions
	<b>Rear-facing Car Seats</b>	Children should use rear-facing car seats in the back seat as long as possible to the rear-facing height and weight limits for the seat (even up to age 2 or 3). If your car seat has a rear-facing weight limit of 22 pounds or less, you should change to a convertible car seat with higher rear-facing limits and keep rear-facing for longer. Leg crowding is expected and okay. It does not cause harm as long as the child is within the weight and height limits for the seat.
	<b>Forward-facing Car Seats with Harnesses</b>	Keep your child rear-facing until the top weight or height limits for the rear-facing car seat. Once top rear-facing limits are reached, use a forward-facing car seat with a harness and a tether. Keep your child in a car seat with a harness until he or she reaches the top height or weight limit for the harness.
	<b>Booster Seats</b>	Use car seats with harnesses to the top weight or height limits for the harnesses. Once children outgrow harnesses, use a booster seat in the back seat <b>until the seat belt fits properly</b> . A booster seat is often needed until a child is around 4 feet 9 inches tall. Your child may be about 12 years old before he/she is ready for a seat belt.
	<b>Seat Belts</b>	Older children should use a lap-shoulder seat belt in the back seat once they outgrow a booster seat. They have not outgrown a booster seat until the seat belt fits correctly: (1) The shoulder strap should cross the center of the chest and rest on the shoulder (not the neck). (2) The lap belt should fit low and snug on the upper thighs (not the stomach). (3) The knees should bend at the edge of the vehicle seat when sitting all the way back.

 [facebook.com/childpassengersafety](https://facebook.com/childpassengersafety)

 [twitter.com/childseatsafety](https://twitter.com/childseatsafety)

[www.nhtsa.gov/Safety/CPS](http://www.nhtsa.gov/Safety/CPS)




Figure 2. Premature Graduation flyer viewed by participants in Group 2

## Right Seat • Right Time • Right Use

Unrestrained children are 3 times more likely to be injured in a crash.

Follow these 4 Steps to Keep Kids Safe in the Car





**1. Keep Kids Rear-Facing as Long as Possible:**  
Children should use rear-facing car seats in the back seat as long as possible to the rear-facing height and weight limits for the seat (even up to age 2 or 3). If your car seat has a rear-facing weight limit of 22 pounds or less, you should change to a convertible car seat with higher rear-facing weight and height limits and keep rear-facing for longer. Leg crowding is expected and okay. It does not cause harm as long as the child is within weight and height limits for the seat.



**2. Keep Kids in Seats with Harnesses as Long as Possible:**  
Once top rear-facing limits are reached, children should use a forward-facing car seat with a harness and tether in the back seat. Use a car seat with a harness as long as possible to the top height or weight limit for the harness.



**3. Use Booster Seats until the Belt Fits:**  
Children should use booster seats in the back seat until the seat belt fits. Kids are not ready for a belt until they pass this test without the booster: (1) They can sit all the way back in the vehicle seat with knees bent at the edge of the seat. (2) The shoulder strap crosses the center of the chest and rests on one shoulder (not the neck). (3) The lap belt fits low & snug on the upper thighs (not the stomach).



**4. Keep Kids Belted in the Back until Age 13:**  
Older children should ride in a lap-shoulder seat belt in the back seat once they outgrow a booster seat, which may not happen until close to 12 years old. The back seat is safest for all children.

 [facebook.com/childpassengersafety](https://facebook.com/childpassengersafety)

 [twitter.com/childseatsafety](https://twitter.com/childseatsafety)

[www.nhtsa.gov/Safety/CPS](http://www.nhtsa.gov/Safety/CPS)



Figure 3. Risk Reduction Rationale flyer viewed by participants in Group 3

## Reducing Car Crash Injury = Right Seat + Right Time + Right Use

- In a crash, the vehicle stops or changes direction in fractions of a second.
- Everyone is thrust in the direction the car was traveling.
- Unrestrained children are 3 times more likely to be injured.
- Your child's restraint is made to stop your child with the vehicle & reduce harm.
- The better the fit to your child's growing body, the better the protection.

**BIRTH**



**HERE'S WHAT TO DO:**

- Start your child in a REAR-FACING CAR SEAT IN THE BACK SEAT. Use as long as possible to the top rear-facing weight and height limits on the seat (even up to age 2 or 3).
- If your car seat has a rear-facing limit of 22 pounds or less, change to a convertible seat with higher rear-facing weight limits. Keep rear-facing longer.
- Leg crowding is expected & okay. It does not cause harm as long as child is within weight & height limits for the seat.

**HERE'S WHY:**

- A rear-facing car seat moves with your child and absorbs crash forces.
- Cradles to reduce harm to neck & spine
- Kids under 2 are more likely to be injured if forward facing.

**HERE'S WHAT TO DO:**

- Once top rear-facing limits are reached, use a FORWARD-FACING CAR SEAT WITH A HARNESS and tether in the back seat.
- Remember to keep rear-facing as long as possible before turning forward.
- Use a car seat with a harness as long as possible to the top height or weight limit for the harness.

**HERE'S WHY:**

- Harnesses spread crash forces over strong parts of the body.
- Keeps body positioned in a crash
- The tether limits head injuries by reducing movement in a crash.

**HERE'S WHAT TO DO:**

- Once a child outgrows the top limits for the harnessed car seat, use a BOOSTER SEAT IN THE BACK SEAT.
- Use a booster seat until the belt fits correctly (see next step).
- A booster seat is often needed until 4 ft 9 in tall. Your child may be 12 years old before ready for a belt.

**HERE'S WHY:**

- A booster raises a child up so the belt rests over strong body parts.
- Decreases stomach, neck, & spine injuries
- Keeps kids from putting the shoulder belt under their arm or behind their back, which causes harm in a crash

**TO TEEN YEARS**



**HERE'S WHAT TO DO:**

- Kids should ride in a lap and shoulder SEAT BELT IN THE BACK SEAT once they outgrow a booster seat.
- Kids are ready for a seat belt when: (1) The shoulder strap crosses the center of the chest and rests on the shoulder (not the neck). (2) The lap belt fits low on the thighs (not the stomach). (3) Knees can bend when sitting all the way back in the seat.
- Use the back seat for kids under 13.

**HERE'S WHY:**

- A seat belt keeps the child in the vehicle.
- Spreads crash forces
- Protects head & spine
- Back seat is safer than the front

**Kids under age 13 are nearly two times safer in the back seat because they are farthest from the most common kind of crash and from frontal airbags. Front airbags are meant for teens and adults.**

 [facebook.com/childpassengersafety](https://facebook.com/childpassengersafety)
 [twitter.com/childseatsafety](https://twitter.com/childseatsafety)
[www.nhtsa.gov/Safety/CPS](http://www.nhtsa.gov/Safety/CPS)


Figure 4. Age-Based flyer viewed by participants in Group 4

## Car Seat Recommendations for Children

- Select a car seat based on your child's age and size, and choose a seat that fits in your vehicle and use it every time.
- Always refer to your specific car seat manufacturer's instructions; read the vehicle owner's manual on how to install the car seat using the seat belt or LATCH system; and check height and weight limits.
- To maximize safety, keep your child in the car seat for as long as possible, as long as the child fits within the manufacturer's height and weight requirements.
- Keep your child in the back seat at least through age 12.



AGE



**Birth – 12 months** 

Your child under age 1 should always ride in a rear-facing car seat. There are different types of rear-facing car seats: Infant-only seats can only be used rear-facing. Convertible and 3-in-1 car seats typically have higher height and weight limits for the rear-facing position, allowing you to keep your child rear-facing for a longer period of time.



**1 – 3 years**  

Keep your child rear-facing as long as possible. It's the best way to keep him or her safe. Your child should remain in a rear-facing car seat until he or she reaches the top height or weight limit allowed by your car seat's manufacturer. Once your child outgrows the rear-facing car seat, your child is ready to travel in a forward-facing car seat with a harness.



**4 – 7 years**  

Keep your child in a forward-facing car seat with a harness until he or she reaches the top height or weight limit allowed by your car seat's manufacturer. Once your child outgrows the forward-facing car seat with a harness, it's time to travel in a booster seat, but still in the back seat.



**8 – 12 years**  

Keep your child in a booster seat until he or she is big enough to fit in a seat belt properly. For a seat belt to fit properly the lap belt must lie snugly across the upper thighs, not the stomach. The shoulder belt should lie snug across the shoulder and chest and not cross the neck or face. Remember: your child should still ride in the back seat because it's safer there.

**DESCRIPTION (RESTRAINT TYPE)**



A **REAR-FACING CAR SEAT** is the best seat for your young child to use. It has a harness and in a crash, cradles and moves with your child to reduce the stress to the child's fragile neck and spinal cord.



A **FORWARD-FACING CAR SEAT** has a harness and tether that limits your child's forward movement during a crash.



A **BOOSTER SEAT** positions the seat belt so that it fits properly over the stronger parts of your child's body.



A **SEAT BELT** should lie across the upper thighs and be snug across the shoulder and chest to restrain the child safely in a crash. It should not rest on the stomach area or across the neck.

 [www.facebook.com/childpassengersafety](https://www.facebook.com/childpassengersafety)
 <http://twitter.com/childseatsafety>
March 21, 2011

Figure 5.  
**Changes in Restraint Selection Score by Flyer Version**




Figure 6.  
**Changes in Rear-facing Knowledge by Flyer Version**




Figure 7.  
**Changes in Self-efficacy by Flyer Version**




Figure 8.  
**Changes in Overall Attitudes by Flyer Version**





Figure 9.  
**Changes in Stated Intentions by Flyer Version**



U.S. Department  
of Transportation  
**National Highway  
Traffic Safety  
Administration**

This and other behavioral-related research notes and reports may be found at [www.nhtsa.gov/BehavioralResearchNotes](http://www.nhtsa.gov/BehavioralResearchNotes).